

TWIGAMEC

PRESENTATION: Twigamec is a clear, colourless, slightly viscous, Sterile solution containing 1% Ivermectin.

Specific Indications: Twigamec is indicated for the control of gastro-intestinal nematodes, lungworms, lice and mange mites.

SPECIES: Cattle, Sheep, Goats and Pigs

DOSAGE & ADMINISTRATION:

Cattle: Dosage: The recommended dose is 1 ml per 50 kg bodyweight.
Administration: Inject subcutaneously in front or behind the shoulder using aseptic technique.

Pigs: Dosage: The recommended dose is 1 ml per 33 kg bodyweight. Administration: Inject subcutaneously into the neck using aseptic technique.

RECOMMENDED WITHHOLDING PERIOD:

Cattle: - Cattle must not be slaughtered for human consumption within 21 days after treatment.
- Do not treat cattle producing milk for human consumption or in dairy cows within 60 days before calving.

Pigs: - Pigs must not be slaughtered for human consumption within 21 days after treatment.

PHARMACEUTICAL PRECAUTIONS:

- Once vial is broached, the product should be used within 28 days.
- The product is extremely dangerous to fish and aquatic life.
- Do not contaminate surface waters or ditches with product container or unused product.

PACKS 50 ml vials.

