

POM

TWIGAMYCIN

Oxytetracycline 10% Injection
For Cattle, Sheep, Goats & Pigs
FOR VETERINARY USE ONLY

WARNING

Keep out of reach of children.
Avoid the introduction of contamination during use.

PACKING

Vial of 50ml and 100ml.



REGISTERED AND DISTRIBUTED BY:
Twiga Chemical Industries Ltd.
P.O. Box 30172 -00100, NAIROBI, KENYA
Tel: +254 20 3942000, Fax: +254 20 3942405



TWIGAMYCIN

Oxytetracycline 10% Injection

For Cattle, Sheep, Goats & Pigs



For
Veterinary
use only

Kwa tiba ya mifugo



REGISTERED AND DISTRIBUTED BY:
Twiga Chemical Industries Ltd.
P.O. Box 30172 -00100, NAIROBI, KENYA
Tel: +254 20 3942000, Fax: +254 20 3942405



POM

TWIGAMYCIN

Oxytetracycline 10% Injection
For Cattle, Sheep, Goats & Pigs

FOR VETERINARY USE ONLY

DESCRIPTION

Oxytetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. The action of oxytetracycline is based on inhibition of bacterial protein synthesis. Oxytetracycline is mainly excreted in urine, for a small part in bile and in lactating animals in milk.

COMPOSITION

Contains per ml.: Oxytetracycline base 100 mg. Solvents ad. 1 ml.

INDICATIONS

Arthritis, gastrointestinal and respiratory infections caused by oxytetracycline sensitive micro-organisms, like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp., in calves, cattle, goats, sheep and swine.

CONTRA-INDICATIONS

- Hypersensitivity to tetracyclines.
- Administration to animals with a serious impaired renal and/or liver function.
- Concurrent administration with penicillines, cephalosporines, quinolones and cycloserine.
- Do not use in cats, dogs and horses.



SIDE EFFECTS

- After intramuscular administration local reactions can occur, which disappear in a few days.
- Discoloration of teeth in young animals.
- Hypersensitivity reactions.

DOSAGE AND ADMINISTRATION

For intramuscular or subcutaneous administration:

Full-grown animals : 1 ml. per 10 - 20 kg. body weight for 3 - 5 days.

Young animals: 2 ml. per 10 - 20 kg. body weight for 3 - 5 days.

Do not administer more than 20 ml. in cattle, more than 10 ml. in swine and more than 5 ml. in calves, goats and sheep per injection site.

WITHDRAWAL TIMES

- For meat : 15 days.
- For milk : 5 days.

STORAGE

Store in a cool dry place away from direct sunlight

Store Below 30°C.

Protect from Light .