

COMPOSITION:

Each ml contains:

Oxytetracycline Hydrochloride Ph. Eur.100mg
Excipient q.s.1ml

CONTRA INDICATIONS WARNINGS, ETC.:

Milk taken from treated cows should not be used for human consumption during treatment and for 60 hours after treatment. Animals must not be slaughtered for human consumption during treatment. Animals may be slaughtered for human consumption only after 15 days from the last treatment.

General toxicity is low although collapse has been reported with tetracycline in weak or debilitated animals.

Manufactured by:

HEBEI YUANZHENG PHARMACEUTICAL
LIABILITY CO. LTD.
SHIJIAZHUANG, HEBEI, CHINA

A product of:

Twiga Chemical Industries Ltd.
P.O. Box 30172 -00100, NAIROBI, KENYA

TWIGAMYCIN

Oxytetracycline 10% Injection in Cattle, Sheep, Goats & Pigs

For Animal Treatment Only

Keep Out of Reach of
Children

Store Below 30°C.

Protect from Light .

Avoid the
introduction of
contamination
during use.

100ML



TWIGA
CHEMICALS

**DOSAGE AND ADMINISTRATION**

By intramuscular or slow intra-venous injection:

Cattle: 1.5 – 4.0 mg per kg body weight daily.
Pigs: – 2.0-9.0 mg per kg body weight daily.

PHARMACEUTICAL PRECAUTIONS:

Store at room temperature, not exceeding 30°C

Protect from light. Keep out of reach of Children.

Following withdrawal of the first dose, use the product within 28 days.
Discard unused material.

MFG:

EXP:

Batch NO.