

DESCRIPTION

Seed dressing fungicide for seed dressing for the control of early seedling, seed and soil borne diseases (*Fusarium sp, Phoma sp, Bipolaris sp, Alternaria alternata*) in wheat.
SEEDGUARD 400FS is a combination of a systemic product, carboxin, which controls diseases within the seed as well as externally, and a general contact fungicide, Thiram, which increases the level of disease control and extends the range of diseases controlled. SEEDGUARD 400FS is recommended for use in the control of soil, seed borne diseases and seedling diseases in wheat.

DIRECTION FOR USE

Disclaimer: Only treat good quality seed as treatment of mechanically damaged or poor quality seed may result in reduced germination, reduced seedling vigor and emergence. Seed to be treated should have moisture content of 15% or less.

Mixing and Application

SEEDGUARD 400FS should be applied as a slurry treatment by means of suitable seed treatment equipment with a uniform distribution chamber. SEEDGUARD 400FS may be applied undiluted. In concrete mixers and machines with only one secondary distribution, SEEDGUARD 400FS may be diluted twice its own volume with clean water. Ensure thorough seed coating by correct calibration of the equipment. Maintain gentle agitation during mixing and application.

Rate of application

Crop	Disease	Rate ML/100kg seed	Comment	Pre-Harvest Interval
Wheat	Seedborne pathogens	300mls	Mix thoroughly and ensure recommended rate is used.	Product is used as a seed dresser
	<i>Alternaria spp</i> (causes kernel smudge, seedling blight, sooty moulds)			
	<i>Bipolaris spp</i> (causes spot blotch)			
	<i>Phoma spp</i> (causes phoma spot)			
	Soil borne pathogens:			
	<i>Fusarium spp</i> (causes root rot, seedling blight, head blight)			

CROP ESTABLISHMENT

At recommended rate, SEEDGUARD 400FS is not phytotoxic to seed and the young establishing plants. This enables seedlings to reach the soil surface quickly and establish strongly

COMPATIBILITY: SEEDGUARD 400FS is compatible with seed dressing insecticides. However it is always recommended that compatibility test be done in a small container before. SEEDGUARD 400FS can be mixed with other products

HAZARD AND PRECAUTION

SEEDGUARD 400FS is harmful if swallowed, inhaled or absorbed through the skin. Eye irritant. Avoid contact with skin, eyes and clothing. Wash hands and face thoroughly with soap and water after use and before eating or smoking. Consumption of alcoholic beverages before and after exposure may enhance the symptoms, which include nausea, vomiting, fatigue or flushing. Remove contaminated clothing immediately after use. Use protective clothing when handling: long sleeved shirts and long pants, water proof gloves, shoes, protective eye wear (face shield) and wash contaminated clothing before re-using. Clearly mark bags containing treated seed "POISONOUS". Empty bags should not be re-used for any other purpose.

Do not use treated seed for feed or food purposes.

STORAGE

Store in original, tightly closed container away from children, food, drinks and livestock feedstuffs. Store in a cool, dry and well ventilated area under lock and key. Keep in a cool dark place away from sparks and flames. Extinguish fire with dry chemicals, foam or water spray. Do not use water jet. Avoid exposure to direct sunlight, heat and fire.

Disposal

Surplus product, if unused should be disposed off on a site approved for pesticide in a safe place away from water sources. Do not use empty container for any other purpose. Empty containers are to be flattened, perforated, and disposed off on a site approved for pesticide disposal according to national regulation on waste disposal.

Spillage and Leakage

Remove contaminated soil layer and dispose off in a safe place away from water sources. Soak up with sand, sawdust, soil or binding agent. Collect the contaminated material thoroughly and dispose off in a safe place, approved for pesticide. Clean spilled area with plenty of water. Spilt product must not be re-used.

ENVIRONMENT AND WILDLIFE

To fish and aquatic invertebrates: Do not contaminate water sources and supplies by disposal of product waste. Keep livestock out of treated areas.

FIRST AID INSTRUCTIONS

If poisoning is suspected, call a physician immediately.

In case of eye contact: Flush the eyes with plenty of water for at least 15 minutes. Contact a physician immediately. **In case of skin contact:** Wash contaminated skin with soap and water. Remove contaminated clothing. If skin irritation develops, contact a physician immediately.

In case of inhalation: Remove the patient away from contaminated area and place the patient in a well-ventilated area. Keep warm and at rest. Consult a doctor.

In case of ingestion: Dilute the swallowed product by giving plenty of water. Do not make the person vomit, as solvents may damage lungs. Take person and this label to the doctor at once.

TOXICOLOGICAL INFORMATION

SEEDGUARD 400FS is harmful if swallowed, inhaled or absorbed through the skin. Symptoms of poisoning: contact with skin or eyes may cause moderate irritation. Ingestion of large amounts together with alcohol may produce nausea, vomiting or diarrhea.

Antidote: No specific antidote is known.

Notes to Physician: Treat symptomatically. No antidote.

In case of poisoning, call Toll free No: 0800720021 or 0800730030 (24hrs)

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY: Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

READ THE LABEL BEFORE USING. (SOMA KIBANDIKO KABLA YA KUTUMIA)

SEED GUARD® 400 FS

Seed dressing fungicide for seed dressing for the control of early seedling, seed and soil borne diseases (*Fusarium sp, Phoma sp, Bipolaris sp, Alternaria alternata*) in wheat.
(Kiukuvu ya kudhibiti magonjwa yanayobebwa kwa mbegu, magonjwa yanayopatikana kwa mchanga na magonjwa ya miche ya ngano)

REGISTRATION NO.: PCPB (CR) 1633
(NAMBARI YA KUSAJILIWA)

GUARANTEE: Carboxin 200g/L +
(DHAMANA) Thiram 200g/L

COMMERCIAL AND AGRICULTURAL CLASS (SEHEMU YA BIASHARA NA KILIMO)

KEEP LOCKED OUT OF REACH OF CHILDREN (WEKA MBALI NA WATOTO)



Treated Wheat Seed

Rotting Wheat Seedlings

Healthy wheat seedlings


Wheat Field

In case of poisoning call Toll free Nos

(Piga simu isiyo na malipo wakati wa kusumika):

0800720021 and 0800730030 (24hrs /masaa 24)

SHELF LIFE: 2 years from date of manufacture if stored in original, unopened container in a cool dry well ventilated place.
(MUHA RAFUNI: Miaka miwili baada ya tarehe ya kutengenezwa ikihifadhiwa kwenye chombo chake asili kischofunguliwa na ilioweka pahali pakavu na penye baridi).



REGISTERED AND DISTRIBUTED IN KENYA BY:

TWIGA CHEMICAL INDUSTRIES LTD.

P.O. BOX 30172, 00100 GPO,

NAIROBI, KENYA

TEL: 3942000 , FAX:- 3942405

DATE OF MANUFACTURE (TAREHE YA KUTENGENEZWA)	March 2018
BATCH NO: NAMBARI YA KIFURUSHI	SNFWAJ1803
EXPIRY DATE (TUMIA KABLA YA):	March 2020

Net Content /Kiasi
1 litre/Lita 1

MANUFACTURER:

Shaanxi Hengtian Chemical Co.Ltd
No. 48 Fengcheng 3 Road, Xi'AN, China

EXPORTER

Ningbo Sunjoy Agrosience Co.Ltd
(Formerly called Sinochem Ningbo Ltd)
21 Jiangxia ST., Ningbo 315000 China

MAELEZO

Kiukuvu ya kudhibiti magonjwa yanayobebwa kwa mbegu, magonjwa yanayopatikana kwa mchanga na magonjwa ya miche ya ngano)

SEEDGUARD 400FS ni mchanganyiko wa dawa inyoilingia ndani ya mmea [systemic]; Carboxin- huzuia magonjwa ndani na miche ya mbegu. Thiram hufanya kazi kwa kugiza, hii huongeza kiwango cha uthibiti wa magonjwa yanayodhibitiwa. SEEDGUARD 400FS inapendekzewa kuzuia magonjwa yanayobebwa kwa mbegu, magonjwa yanayopatikana kwa mchanga na magonjwa ya miche ya ngano.

MAAGIZO KWA MATUMIZI

Onyo: Tibu tu mbegu nzuri. Ukutibu mbegu duni inaweza kusababisha kupungua kuota , kupunguza miche kuibuka . Wakati wa kutibu Mbegu nafaa kuwa na unyevu wa 15% au chini.

Mchanganyika na jinsi ya kutumia

SEEDGUARD 400FS inapaswa kutumika kama tiba tope kwa njia ya kufaa kwa vifaa vya kutibu mbegu na chumba msingi ya usambazaji. SEEDGUARD 400FS inaweza kutumika bila kunyanyagaa kwa maji. Katika mitambo ya kunchanganya simiti na katika mashini ambapo usambazaji hufanywa kama hatua ya pili. SEEDGUARD 400FS inaweza kuchanganywa kwa kiasi cha hadi mara mbili yake kwa maji. Hakikisha kwamba vifaa ama mashini ya kuweka dawa zimesetiwa vizuri ili mbegu zikolee dawa vizuri. Endelea kukoroga polepole wakati wa kuchanganya na kupaka.

VIWANGO YAA KUTUMIA

Mimea	Magonjwa	Kiwango wa dawa ML/100kg wa Mbegu	Muda kabla ya kuvunja	PHI
Ngano	Magonjwa yanayobebwa kwa mbegu	Mililita 300	Changanya vizuri, hakikisha kiwango cha dawa kinachofaa kinatumika	Dawa hii hutumika kwa kuchanganya na mbugu kabla ya mbegu kupandwa
	<i>Alternaria spp</i> (causes kernel smudge, seedling blight, sooty moulds)			
	<i>Bipolaris spp</i> (causes spot blotch)			
	<i>Phoma spp</i> (causes phoma spot)			
	Magonjwa yanayopatikana kwa mchanga			
	<i>Fusarium spp</i> (causes root rot, seedling blight, head blight)			

KUANZISHWA KWA MAZAO

Kwa kiwango inavyopendekzewa, SEEDGUARD 400FS haiwezi kuimiza miche, Hii itawezesha miche kumea kwa haraka na kuanzisha kukaa vizuri.

UPATANFU: SEEDGUARD 400FS

hupatana na dawa za kuangamiza wadudu zinazochanganywa na mbegu. Lakini ni muhimu kufanya majaribio yako binafsi kuhakikisha kuwa haina mahadhara katika hali yako, kabla SEEDGUARD 400FS ichanganywe na madawa zingine.

HATARI NA TAHADHARI

SEEDGUARD 400FS ni hatari kimezwa, ikihutwa ndani au ikinyonywa ndani ya ngozi. Inawasha macho, Isiguze ngozi, macho au ngozi. Osha mikono na uso vizuri kwa sabuni na maji baada ya kutumia na kabla ya kula chakula au kuvuta sigara. Unyuaji wa pombe kabla na baada ya kutumia dawa hii huenza ukazidisha dalili kama za kuchafuka moyo, kutapika au uchovu. Toa ngozi zilichogwa dawa mara moja baada ya kazi. Tumia mavazi ya ya kujikinga unapo tumia dawa hii kama mashaji ya mikono mrefu na sunati ndetu govu za mpira vya kinga ya macho na uoshie nguo uliroyanika kazi kabla ya kutumia tena. Andika wazi neno 'SUMU' kwenye mikoto yote iliyo na mbegu zilizowekea dawa, ili hii isitumike tena. Usitumie mbegu zilizowekea dawa kama chakula au malisho.

KUHIFADHI

Hifadhi ndani ya chombo chake asili kischofunguwa vizuri mbali na watoto, chakula, vinywaji na chakula cha mifugo. Hifadhi SEEDGUARD 400FS mahali baridi, pakavu na panapoitikisha hewa safi na paipofunguwa na kufufu.

KUTUPA

Dawa ya ziada kama hakikutumia itupwa kwenye jaa lililohinshwa ili kutupa madawa ya sumu ya kuangamiza wadudu waharibu mahali salama mbali na vyanzo vya maji. Usitumie chombo kilichokwisha dawa kwa matumizi mengine yeyote.

Kimwagika au kifuja

Ondoa mchanga wa juu uliionga dawa na utupe pahali salama mbali vyanzo vya maji. Kimwagika inonywe kwa kuimwagika mchanga, unga wa mibao udogo wa mibao kumasa. Kusanya kwa makini vifu vilivyio njia dawa na utupe pahali salama lililohinshwa. Safisha kwa maji eneo lililomwagika na dawa. Dawa iliyomwagika istumike tena.

MADHARA KWA MAZINGIRA

SEEDGUARD 400FS ni sumu kwa samaki, usichafue mto, mabwawa, visime na vyanzo au njia za maji kwa kusafisha vifaa vilivyotumika, ama kutupa masaa zote.

MAAGIZO YA HUDUMA YA KWANZA

Kama unashuku sumu, ita daktari mara moja.

Kiwa imeingia ndani ya macho: Osha macho na maji mengi kwa angalau dakika 15. Wasiliana na daktari mara moja.

Kiwa imeguza ngozi: Osha ngozi ili ingie dawa kwa sabuni na maji. Toa ngozi zilimwagika na dawa.

Kiwa mwasho wa ngozi utokeze, onana na daktari mara moja.

Kiwa mvuke wa dawa hii unevutwa ndani ya mapafu: Muondoe mgongo kutoka kwenye eneo ya sumu na umweke mgonjwa mahali penye hewa safi. Mweke kwenye joto la kadiri na katika hali ya kupumzikia. Pata ushaua wa daktari.

Kama ime meza: Ipinguze nguvu ya dawa iliyomezwa haraka kwa kunnywesha mgonjwa maji mengi usifanye atapike. Muone daktari haraka.

MAELEZO YA SUMU

SEEDGUARD 400FS ni hatari kimezwa au ikilinga kwa mwili kupitia kwa mapua au ngozi.

Dalili za za sumu ikiitumika kwenye mgonjwa inaweza kusababisha mwasho wa kadiri. Kimezwa kwa wingi pamoja na pombe inaweza kusababisha kuchafuka moyo, kutapika ama kuhara.

Dawa ya kuvunja nguvu za sumu: Hakuna dawa maalum ya kuvunja nguvu za dawa hii.

Maelezo kwa daktari

Tibu kidhana na dalili

Piga simu isiyo na malipo wakati wa kusumika: 0800720021 au 0800730030 (masaa 24)

ILANI KWA MATUMIAJI: Dawa hii sharti itumie kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya Sheria za Madawa ya kuangamiza wadudu waharibu kutumia na kuhifadhi dawa hii katika hali yote salama.

THIBITISHO: THIBITISHO: Hii ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu, Munuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binaadamu au maji na lazima akubali dawa hii katika hali hiyo.